

On $N = 2$ M QCD [†]

Toshio Nakatsu

*Department of Physics,
Graduate School of Science, Osaka University,
Toyonaka, Osaka 560, JAPAN*

Abstract

We review M -theory description of $4d$ $N = 2$ SQCD. Configurations of M -theory fivebranes relevant to describe the moduli spaces of the Coulomb and Higgs branches are studied using the Taub-NUT geometry. Minimal area membranes related with the BPS states of $N = 2$ SQCD are given explicitly. They almost saturate the BPS bounds. The deviation from the bounds is due to their boundary condition constrained by the fivebrane. The electric-magnetic duality at the baryonic branch root is also examined from the M -theory viewpoint. In this course, novel concepts such as creation of brane and exchange of branes in Type II theory are explained in the framework of M -theory.

[†] Lecture in the Second Winter School on “Branes, Fields And Mathematical Physics” at the APCTP (Feb.9-20, 1998).

Many interesting results about gauge field theories in various dimensions have been obtained by analyzing the worldvolume effective theory of branes in superstring theory [1]. They can be realized by branes mostly in Type IIA or IIB superstring theory, but particularly an interesting configuration, which describes four-dimensional $N=2$ supersymmetric QCD ($4d\ N=2$ SQCD), has been proposed in [2] within the framework of M -theory. In this construction a mysterious hyper-elliptic curve, which is used for the description of the exact solution of the Coulomb branch of $4d\ N=2$ SQCD, appears as a part of a M -theory fivebrane.

It is pointed out [3, 4, 5, 6] that there exist various dualities between supersymmetric gauge field theories and that these dualities play an important role for our understanding of their non-perturbative dynamics. Several steps to clarify an origin of these dualities from the string theory viewpoint have been taken [7, 8]. In this course of explanation we need novel concepts such as a brane can be created or annihilated when two different branes cross each other. However, since these phenomena are due to the strong coupling dynamics of Type II theories, it is still difficult to treat them correctly in these theories. On the other hand, M -theory includes the strong coupling dynamics of Type IIA theory in its semi-classical description [9]. So one can expect that the “dualities” accompanied by exchanging branes can be understood via semi-classical analysis of M -theory.

In this article we review M -theory description of $4d\ N=2$ SQCD. Configurations of M -theory fivebranes relevant to describe the moduli spaces of the Coulomb and Higgs branches are studied using the Taub-NUT geometry. Minimal area membranes related with the BPS states of $N=2$ SQCD are given explicitly, but, due to their boundary condition, they can not saturate the BPS bounds at a generic point of the moduli space of the configuration. This signals a necessity to incorporate some recoil of the membranes to the fivebrane. The electric-magnetic duality at the baryonic branch root of $4d\ N=2$ SQCD is also examined from the M -theory viewpoint. In this course, novel concepts such as creation of brane and exchange of branes in Type II theory are explained in the framework of M -theory.

The organization is as follows : In section 1 we start with a realization of the Coulomb branch of $4d\ N=2$ SQCD by M -theory fivebrane configuration. In this description the worldvolume of fivebrane includes the so-called Seiberg-Witten curve, as a part. An inclusion of matters is expressed by an embedding of the curve into the multi-Taub-NUT space. This embedding is studied in detail by using the concrete metric of the multi-Taub-NUT space. In section 2 minimal area membranes are studied treating the previous M -theory brane configuration as their background geometry. These membranes are classified by their shapes in ten-dimensions

and boundary conditions in eleven-dimensions. In section 3 we study the Higgs branch of $4d$ $N = 2$ SQCD. An exact description of the baryonic and non-baryonic branches in terms of M -theory is presented. The s -rule [7] is also derived using the Taub-NUT geometry. In the last section the baryonic branch root, that is, intersection of the Coulomb branch and the baryonic branch is examined from the M -theory viewpoint. It is known [10] in field theory that the baryonic branch root admits to have two descriptions dual to each other. After providing M -theory proof of “brane creation”, we present a specific family of M -theory configurations relevant to explain this duality and examine the BPS states of membranes associated with them.

1 Coulomb Branch of $N=2$ MQCD

Four-dimensional gauge theories with $N = 2$ supersymmetry can be realized [2] as effective theories on the world-volume of a M -theory fivebrane. A different type of gauge theories requires a different topology of M -theory fivebrane and a different eleven-dimensional background where the fivebrane is embedded. This realization of $N=2$ supersymmetric QCD via the world-volume effective theory of the M -theory fivebrane is called $N=2$ M -theory QCD. (MQCD for short.) MQCD does not exactly coincide with an ordinary supersymmetric QCD in four-dimension, but is considered to belong to the same universality class. Moreover many difficulties appearing in the field theoretical analysis of the supersymmetric QCD vacua, which are mainly due to their singularities, are resolved within the framework of M -theory. So, MQCD is a very useful tool for our understanding of the dynamics of supersymmetric QCD.

Consider an eleven-dimensional manifold $M^{1,10}$ of M -theory which admits to have the form

$$M^{1,10} \simeq \mathbf{R}^{1,3} \times X^7. \quad (1.1)$$

$\mathbf{R}^{1,3}$ is the four-dimensional space-time where $N = 2$ supersymmetric theory will exist. X^7 is a (non-compact) seven-dimensional manifold which suffers several constraints due to the requirement of $N = 2$ supersymmetry on the worldvolume. The supersymmetry of M -theory in the eleven-dimensions is generally broken by this product space structure of $M^{1,10}$. However, if the submanifold X^7 admits to have a holonomy group smaller than $SO(7)$, some of supersymmetries will survive on the four-dimensional space-time $\mathbf{R}^{1,3}$. Recall that we ultimately realize $N = 2$ supersymmetry on the worldvolume of a M -theory fivebrane, strictly speaking, on its four-dimensional part which is identified with $\mathbf{R}^{1,3}$ in (1.1). The fivebrane itself will be intro-

duced soon later as a BPS saturated state which breaks half the surviving supersymmetries. So, with this reason, we must take X^7 as a submanifold which keeps $N = 4$ supersymmetry on the four-dimensional space-time $\mathbf{R}^{1,3}$. The holonomy group of X^7 is required to be $SU(2)$. This requirement reduces the seven-manifold X^7 to be

$$X^7 \simeq \mathbf{R}^3 \times Q^4, \quad (1.2)$$

where Q^4 is a four-manifold with $SU(2)$ holonomy, that is, a hyper-Kähler manifold. This hyper-Kähler manifold should be chosen appropriately according to whether the theory on $\mathbf{R}^{1,3}$ contains matter hypermultiplets or not.

1.1 Pure $N=2$ MQCD

Let us describe a configuration of M -theory fivebrane suitable to our purpose. Since we want to leave $N = 2$ supersymmetry on $\mathbf{R}^{1,3}$ as the supersymmetry of the worldvolume effective theory of fivebrane, the worldvolume itself must fill all of $\mathbf{R}^{1,3}$. The rest of the fivebrane is a two-dimensional surface Σ in X^7 . The Lorentz group $SO(3)$ of $\mathbf{R}^3$ in (1.2) turns out to be the $SU(2)_R$ -symmetry of $N = 2$ supersymmetry algebra in four-dimensions. In order to preserve this symmetry Σ must lie at a single point in $\mathbf{R}^3$. It can only spread in Q^4 as a two-dimensional surface. To summarize, the worldvolume of the fivebrane must be $\mathbf{R}^{1,3} \times \Sigma$ where $\mathbf{R}^{1,3}$ is identified with the four-dimensional space-time and Σ is a two-dimensional surface embedded in Q^4 .

Further restriction on the fivebrane world-volume is that it must be BPS-saturated in order to preserve a half of supersymmetries. This is achieved by the minimal area embedding of Σ into Q^4 . The area A_Σ of the surface Σ is bounded from the below

$$A_\Sigma \geq \int_\Sigma \Omega_{\mathbf{R}}, \quad (1.3)$$

where $\Omega_{\mathbf{R}}$ is the Kähler form of the hyper-Kähler manifold Q^4 . This inequality is saturated if and only if Σ is holomorphically embedded in Q^4 . This means that, if one introduces the holomorphic coordinates y and v of Q^4 , the surface Σ is a curve in Q^4 defined by a holomorphic function

$$F(y, v) = 0. \quad (1.4)$$

Let Σ be a Riemann surface with genus $N_c - 1$. Then there appear $N_c - 1$ massless vector multiplets on $\mathbf{R}^{1,3}$ [11]. So, the effective theory on $\mathbf{R}^{1,3}$ is a supersymmetric $U(1)^{N_c-1}$ gauge

theory. In particular let Σ be the Seiberg-Witten curve Σ_{SW} [3, 4] for the pure $SU(N_c)$ gauge theory [12, 13].

$$y^2 - 2 \prod_{a=1}^{N_c} (v - \phi_a) y + \Lambda^{2N_c} = 0. \quad (1.5)$$

We now take, as the four-manifold Q^4 , a flat space $Q_0 \equiv \mathbf{R}^3 \times S^1$ with coordinates (x^4, x^5, x^6, x^{10}) . S^1 is the circle of the eleven-dimension with radius R . For the later convenience we rescale the coordinates and define

$$(v, b, \sigma) \equiv \frac{2}{R}(x^4 + ix^5, x^6, x^{10}). \quad (1.6)$$

The complex structure appropriate to describe the holomorphic embedding of the Seiberg-Witten curve is given [2] by v and $y(b, \sigma) = e^{-\frac{b+i\sigma}{2}}$. Using these holomorphic coordinates the (trivial) hyper-Kähler structure of Q_0 can be written as

$$\begin{aligned} \Omega_{\mathbf{R}} &= \frac{iR^2}{2} \left(\frac{1}{4} dv \wedge d\bar{v} + \frac{dy}{y} \wedge \frac{\overline{dy}}{y} \right), \\ \Omega_{\mathbf{C}} &= \frac{iR^2}{2} dv \wedge \frac{dy}{y}, \end{aligned} \quad (1.7)$$

where $\Omega_{\mathbf{R}, \mathbf{C}}$ are respectively Kähler and holomorphic two-forms ¹.

Let us comment on the holomorphic embedding of Σ_{SW} into Q_0 . One can visualize it by considering eq.(1.5) in terms of v, b and σ , that is, considering it in the real coordinates of Q_0 . Recall the curve is hyper-elliptic. Let $\Sigma_{\pm}$ be its two Riemann sheets

$$\Sigma_{SW} = \Sigma_+ \cup \Sigma_-. \quad (1.8)$$

These Riemann sheets $\Sigma_{\pm}$ are embedded into Q_0 by the equations

$$y_{\pm}(v) = y(b, \sigma), \quad (1.9)$$

where $y_{\pm}(v)$ are two roots of eq.(1.5)

$$y_{\pm}(v) = \prod_{a=1}^{N_c} (v - \phi_a) \pm \sqrt{\prod_{a=1}^{N_c} (v - \phi_a)^2 - \Lambda^{2N_c}}, \quad (1.10)$$

and the holomorphic coordinate y is regarded as the function of b and σ .

¹We normalize $\Omega_{\mathbf{R}, \mathbf{C}}$ so that $dvol_{Q_0} = \frac{1}{2} \Omega_{\mathbf{R}} \wedge \Omega_{\mathbf{R}} = \frac{1}{4} \Omega_{\mathbf{C}} \wedge \overline{\Omega}_{\mathbf{C}}$, where $dvol_{Q_0}$ is the volume form of Q_0 , $dx^4 \wedge dx^5 \wedge dx^6 \wedge dx^{10}$.




Figure 1: *M*-theory fivebrane and Type IIA configuration.

These Riemann sheets are patched together in Q_0 by N_c one-dimensional circles. Let us denote them C_a ($1 \leq a \leq N_c$). It holds

$$\Sigma_+ \cap \Sigma_- = C_1 \cup \cdots \cup C_{N_c} \quad (\subset Q_0). \quad (1.11)$$

Each C_a can be regarded as the S^1 of Q_0 , that is, the circle of eleven-dimension. While this, if considering it as a one-dimensional cycle of the curve, C_a is a cut of the Riemann sheets, that is, the α -cycle of the curve.

These N_c circles on Σ_{SW} or presumably their infinitesimally neighborhoods in Σ_{SW} become N_c D fourbranes in Type IIA theory. The residual part of the curve becomes two NS fivebranes. After all, the single fivebrane in *M*-theory described by eq.(1.5) becomes two NS fivebranes with worldvolume $(x^0, x^1, x^2, x^3, x^4, x^5)$ and N_c D fourbranes with worldvolume $(x^0, x^1, x^2, x^3, [x^6])^2$ stretching between them in the x^6 -direction. See Fig.1.

1.2 $N=2$ MQCD with matter

In Type IIA picture an inclusion of N_f matter hypermultiplets into the above pure gauge theory can be achieved by considering N_f D sixbranes with worldvolume $(x^0, x^1, x^2, x^3, x^7, x^8, x^9)$ and then putting these sixbranes between two NS fivebranes. In such a configuration there appears $N=2$ $SU(N_c)$ supersymmetric QCD with N_f flavors on their common worldvolume (x^0, x^1, x^2, x^3) . This is because the open string sector between N_c D fourbranes and N_f D sixbranes of the configuration includes hypermultiplets which belong to the fundamental representations both of the gauge and flavor groups.

² $[x^6]$ describes a finite interval.

1.2.1 Sixbranes and the multi-Taub-NUT space

Sixbranes can be regarded as the “Kaluza-Klein monopoles” of M -theory compactified to Type IIA theory with a circle S^1 . This is because they are magnetically charged with respect to the $U(1)$ gauge field associated with this S^1 .

Consider a single sixbrane solution of Type IIA SUGRA. Let us denote the worldvolume and transverse directions of a sixbrane respectively by $\vec{x}_{\parallel}$ and $\vec{x}_{\perp}$ ³. The solution [14] can be written by using a single potential V

$$V(x_{\perp}) = 1 + \frac{g_s l_s / 2}{x_{\perp}}, \quad (1.12)$$

where $x_{\perp} \equiv \sqrt{\vec{x}_{\perp} \cdot \vec{x}_{\perp}}$. The ten-dimensional metric and the dilaton have the forms

$$\begin{aligned} ds_{10}^2 &= V(x_{\perp})^{-\frac{1}{2}} d\vec{x}_{\parallel} \cdot d\vec{x}_{\parallel} + V(x_{\perp})^{\frac{1}{2}} d\vec{x}_{\perp} \cdot d\vec{x}_{\perp}, \\ e^{\phi} &= V(x_{\perp})^{-\frac{3}{4}}. \end{aligned} \quad (1.13)$$

The $U(1)$ gauge field A , which has components only in the transverse directions, is determined by⁴

$$\vec{\nabla}_{x_{\perp}} \times \vec{A} = \vec{\nabla}_{x_{\perp}} V(x_{\perp}). \quad (1.14)$$

Inserting them to $ds_{11}^2 = e^{-\frac{2}{3}\phi} ds_{10}^2 + e^{\frac{4}{3}\phi} (dx^{10} + \vec{A} \cdot d\vec{x}_{\perp})^2$, we obtain the eleven-dimensional solution :

$$ds_{11}^2 = d\vec{x}_{\parallel} \cdot d\vec{x}_{\parallel} + \left\{ V(x_{\perp}) d\vec{x}_{\perp} \cdot d\vec{x}_{\perp} + \frac{1}{V(x_{\perp})} (dx^{10} + \vec{A} \cdot d\vec{x}_{\perp})^2 \right\}, \quad (1.15)$$

where the circle of the eleven-dimension has radius $R = g_s l_s$. ($x^{10} \sim x^{10} + 2\pi R$.) The transversal four-dimensional part of this metric describes [15] the Taub-NUT space. To make it transparent we rescale the transversal four-dimensional coordinates to

$$(\vec{r}, \sigma) \equiv \frac{2}{R} (\vec{x}_{\perp}, x^{10}). \quad (1.16)$$

With these rescaled coordinates the eleven-dimensional metric (1.15) becomes

$$ds_{11}^2 = d\vec{x}_{\parallel} \cdot d\vec{x}_{\parallel} + R^2 \left\{ \frac{1}{4} \left(1 + \frac{1}{r} \right) d\vec{r} \cdot d\vec{r} + \frac{1}{4 \left(1 + \frac{1}{r} \right)} (d\sigma + \vec{\omega} \cdot d\vec{r})^2 \right\}, \quad (1.17)$$

³In this article $\vec{x}_{\parallel} = (x^0, x^1, x^2, x^3, x^7, x^8, x^9)$ and $\vec{x}_{\perp} = (x^4, x^5, x^6)$.

⁴ $\vec{\nabla}_{x_{\perp}} \equiv (\partial_{x^4}, \partial_{x^5}, \partial_{x^6})$.

where the transversal part acquires the standard form [16] of the Taub-NUT metric (scaled by R^2).

When there exist N_f parallel sixbranes their transversal four-dimensions becomes the multi-Taub-NUT space. The multi-Taub-NUT metric acquires the following standard form [16] in our coordinates :

$$ds^2 = R^2 \left\{ \frac{V(\vec{r})}{4} d\vec{r} \cdot d\vec{r} + \frac{1}{4V(\vec{r})} (d\sigma + \vec{\omega} \cdot d\vec{r})^2 \right\}, \quad (1.18)$$

The potential $V(\vec{r})$ is given by the superposition

$$V(\vec{r}) = 1 + \sum_{i=1}^{N_f} \frac{1}{|\vec{r} - \vec{r}_i|}, \quad (1.19)$$

where $\vec{r}_i$ denotes the position of the i -th sixbrane. The $U(1)$ gauge field $A = \vec{\omega} \cdot d\vec{r}$ is determined by the relation

$$\vec{\nabla}_r \times \vec{\omega} = \vec{\nabla}_r V(\vec{r}). \quad (1.20)$$

In order to proceed further let us give some comments on sixbranes and the Taub-NUT geometry. First, the positions of sixbranes denoted by $\vec{r}_i$ look like singularities in the metric (1.18). This is incorrect. Actually, parallel sixbranes are the removable NUT singularities in their transversal multi-Taub-NUT space as far as they do not coincide. This means that *sixbranes become geometry in M-theory*. Second, the multi-Taub-NUT space is known to be a hyper-Kähler manifold. Regarding it as a complex manifold N_f parallel sixbranes play the role of resolution or deformation of the simple singularity of type A_{N_f-1} .

1.2.2 Hyper-Kähler structure of the multi-Taub-NUT space

We want to consider a configuration of M -theory fivebrane which realize $N = 2$ SQCD with N_f flavors. For this purpose two-dimensional part Σ of the fivebrane must be embedded into the above multi-Taub-NUT space. To discuss its BPS saturation we need the hyper-Kähler structure [17] [18] of the multi-Taub-NUT space.

Let us introduce the complex structure [17] [18] of the multi-Taub-NUT space Q appropriate to describe the BPS saturation of the fivebrane. For this purpose we need to comment on the Dirac strings associated with the sixbranes. If one regards Q as a Riemannian manifold the direction of the Dirac strings is arbitrary. But, once one introduces a complex structure, their direction becomes unique. This is because putting a complex structure means restricting the

structure group from $SO(4)$ to $U(2)$. This $U(2)$ simply acts as $SO(2)$ on $\vec{r}$ and can not change the direction of the Dirac strings. So, taking it reverse, in order to introduce a complex structure we need to fix the Dirac strings of the sixbranes. Let us separate $\vec{r}$ into two parts $\vec{r} = (v, b)$. (See eq.(1.6).) Denote the position of the i -th sixbrane by $\vec{r}_i = (e_i, b_i)$. We will fix the i -th Dirac string as the semi-infinite line in the (v, b) -space running from $\vec{r}_i = (e_i, b_i)$, parallel with the b -axis, into $(e_i, +\infty)$. They are associated with the following $U(1)$ gauge field $A = \vec{\omega} \cdot d\vec{r}$:

$$\vec{\omega} \cdot d\vec{r} = \text{Im} \left(\sum_{j=1}^{N_f} \frac{1}{\Delta_j} \frac{b - b_j + \Delta_j}{v - e_j} dv \right), \quad (1.21)$$

where

$$\begin{aligned} \Delta_j &\equiv \sqrt{(b - b_j)^2 + |v - e_j|^2} \\ &= |\vec{r} - \vec{r}_j|. \end{aligned} \quad (1.22)$$

The field strength of A becomes

$$F \sim \pi i \sum_{j=1}^{N_f} \theta(b - b_j) \delta^{(2)}(v - e_j) dv \wedge \overline{dv}, \quad (1.23)$$

which shows that the Dirac strings are fixed in the way we expect.

Now let us introduce the following function $y(v, b, \sigma)$ on Q :

$$y(v, b, \sigma) \equiv C e^{-\frac{b+i\sigma}{2}} \prod_{j=1}^{N_f} (-b + b_j + \Delta_j)^{\frac{1}{2}}, \quad (1.24)$$

where C is some constant. Notice that the differential dy can be expressed in terms of dv, db and $d\sigma$:

$$\frac{2dy}{y} = -V db - i d\sigma + \text{Re}(\delta dv), \quad (1.25)$$

where we abbreviate the notation

$$\delta \equiv \sum_{i=1}^{N_f} \frac{1}{\Delta_i} \frac{b - b_i + \Delta_i}{v - e_i}. \quad (1.26)$$

The multi-Taub-NUT metric (1.18) with the fixed gauge potential (1.21) can be rewritten only in terms of v and y :

$$ds^2 = R^2 \left[\frac{V}{4} (db^2 + dv d\bar{v}) + \frac{1}{4V} \{d\sigma + \text{Im}(\delta dv)\}^2 \right]$$

$$\begin{aligned}
&= R^2 \left[\frac{V}{4} dv d\bar{v} + \frac{1}{4V} \left\{ V^2 db^2 + (d\sigma + \text{Im}(\delta dv))^2 \right\} \right] \\
&= R^2 \left[\frac{V}{4} dv d\bar{v} + \frac{1}{4V} \left\{ \{ \text{Re}(2dy/y - \delta dv) \}^2 + \{ \text{Im}(2dy/y - \delta dv) \}^2 \right\} \right] \\
&= R^2 \left\{ \frac{V}{4} dv d\bar{v} + \frac{1}{4V} \left(\frac{2dy}{y} - \delta dv \right) \overline{\left(\frac{2dy}{y} - \delta dv \right)} \right\}, \tag{1.27}
\end{aligned}$$

where we utilize eq.(1.25) to derive the third equality. This shows that y in (1.24) and v give the holomorphic coordinates of Q with which the metric becomes Kählerian. The Kähler and holomorphic two-forms are given by

$$\begin{aligned}
\Omega_{\mathbf{R}} &= \frac{iR^2}{2} \left\{ \frac{V}{4} dv \wedge d\bar{v} + \frac{1}{4V} \left(\frac{2dy}{y} - \delta dv \right) \wedge \overline{\left(\frac{2dy}{y} - \delta dv \right)} \right\}, \\
\Omega_{\mathbf{C}} &= \frac{iR^2}{2} dv \wedge \frac{dy}{y}. \tag{1.28}
\end{aligned}$$

1.2.3 Embedding of the Seiberg-Witten curve

Since the sixbranes play the role of matter hypermultiplets, the Seiberg-Witten curve which is a part of the fivebrane changes [2] to the curve of $N=2$ supersymmetric QCD with N_f flavors [19, 20]

$$y^2 - 2 \prod_{a=1}^{N_c} (v - \phi_a) y + \Lambda^{2N_c - N_f} \prod_{i=1}^{N_f} (v - e_i) = 0, \tag{1.29}$$

where e_i are interpreted as the bare masses of matter hypermultiplets and identified with the positions of sixbranes in the v -plane. Now the above hyper-elliptic curve Σ_{SW} is embedded into the multi Taub-NUT space Q . Let $\Sigma_{\pm}$ be its two Riemann sheets

$$\Sigma_{SW} = \Sigma_+ \cup \Sigma_-. \tag{1.30}$$

These Riemann sheets $\Sigma_{\pm}$ are realized in Q by the equations

$$y_{\pm}(v) = y(v, b, \sigma), \tag{1.31}$$

where $y_{\pm}(v)$ are two roots of eq.(1.29)

$$y_{\pm}(v) = \prod_{a=1}^{N_c} (v - \phi_a) \pm \sqrt{\prod_{a=1}^{N_c} (v - \phi_a)^2 - \Lambda^{2N_c - N_f} \prod_{i=1}^{N_f} (v - e_i)}, \tag{1.32}$$

and the holomorphic coordinate y is regarded as the function (1.24) of v, b and σ .

2 Minimal area membranes and BPS states

In this section we study minimal area membranes which are related with the BPS states of $N = 2$ SQCD. M -theory brane configuration described in the previous section is treated as background geometry for these membranes.

Consider a membrane which worldvolume is $\mathbf{R} \times D$, where D is a two-dimensional surface embedded in the multi-Taub-NUT space Q . Area of the membrane measured by the induced metric satisfies the inequality [21, 22, 23]

$$A_D \geq \left| \int_D \Omega_{\mathbf{C}} \right|. \quad (2.1)$$

The equality, that is, the BPS saturation occurs [22] when D satisfies the following two conditions

- $\Omega_{\mathbf{R}}$ vanishes on D .
- $\Omega_{\mathbf{C}}$ has a constant phase on D .

These membranes will have their correspondences in Type IIA theory. Let π be the following natural projection

$$\pi: Q \longrightarrow \mathbf{R}^3 \quad \pi(v, b, \sigma) = (v, b). \quad (2.2)$$

It is simply a map forgetting about the eleven-dimensional circle of M -theory. When $\dim_{\mathbf{R}} \pi(D) = 1$, the membrane will give rise to a fundamental string in Type IIA theory. When $\dim_{\mathbf{R}} \pi(D) = 2$, It will give rise to a D twobrane.

2.1 Fundamental strings

In this case $\dim_{\mathbf{R}} \pi(D) = 1$ holds. Let D be a two-dimensional cylinder. The boundary of D consists of two circles. $\partial D = S^1 \cup S^1$. The membrane will be classified by its boundary condition in Q . There are three cases :

1. Both two boundary circles of D are on Σ_{SW} in Q and, regarding them as one-dimensional cycles of Σ_{SW} , are homotopic to the α -cycles.
2. One boundary circle of D is on Σ_{SW} in Q and, regarding it a one-dimensional cycle of Σ_{SW} , is homotopic to the α -cycle. The other boundary circle is attached at the sixbrane.

3. Both two boundary circles of D are attached at the sixbranes.

Membrane satisfying the first boundary condition becomes a 4-4 string in Type IIA theory and provides a vector multiplet of the worldvolume gauge theory. Membrane satisfying the second boundary condition becomes a 4-6 string in Type IIA theory and provides a hypermultiplet of the worldvolume gauge theory. Membrane with the last boundary condition becomes a 6-6 string in Type IIA theory and is irrelevant in our discussion because of the decoupling from the fivebrane.

Before presenting investigations on these boundary conditions we examine an implication of the condition, $\dim_{\mathbf{R}}\pi(D)=1$. Let us write down the symplectic forms $\Omega_{\mathbf{R},\mathbf{C}}$ in terms of v, b and σ using the relation (1.25) :

$$\begin{aligned}\Omega_{\mathbf{R}} &= \frac{iR^2}{4} \left\{ \frac{V}{2} dv \wedge d\bar{v} - i db \wedge (d\sigma + \text{Im}(\delta dv)) \right\}, \\ \Omega_{\mathbf{C}} &= \frac{iR^2}{4} dv \wedge \left(\frac{\bar{\delta}}{2} d\bar{v} - V db - i d\sigma \right).\end{aligned}\tag{2.3}$$

Since $\pi(D)$ is a one-dimensional object in the $\mathbf{R}^3$, $dv \wedge d\bar{v}$ and $db \wedge dv$ will vanish on D . It means that the symplectic forms restricted on D have the forms

$$\Omega_{\mathbf{R}}|_D = \frac{R^2}{4} db \wedge d\sigma \quad , \quad \Omega_{\mathbf{C}}|_D = \frac{R^2}{4} dv \wedge d\sigma.\tag{2.4}$$

While this, since D is a two-dimensional cylinder, $d\sigma$ can not vanish on D . So, the vanishing of $\Omega_{\mathbf{R}}$ on D , which one needs for the BPS saturation, requires that *D is included in the hypersurface of Q characterized by $b=\text{const}$.*

2.1.1 4-4 strings

Let us consider a membrane which satisfies the first boundary condition. Namely ∂D is on Σ_{SW} in Q and are homotopic to the α -cycles in Σ_{SW} . Since $\pi(D)$ is one-dimensional, the projection of each boundary circle of D can be expected to be a point on $\pi(\Sigma_{SW})$ or a one-dimensional line on $\pi(\Sigma_{SW})$. What actually occurs is the latter. To explain this, let us recall two Riemann sheets $\Sigma_{\pm}$ of Σ_{SW} are patched together in Q by N_c circles C_a :

$$\Sigma_+ \cap \Sigma_- = C_1 \cup \cdots \cup C_{N_c} \quad (\subset Q).\tag{2.5}$$

These circles, if one regards them as one-dimensional cycles of Σ_{SW} , are homotopic to the α -cycles. Notice that they are projected by π to one-dimensional lines L_a in the $\mathbf{R}^3$

$$L_a = \pi(C_a) \quad (\subset \mathbf{R}^3),\tag{2.6}$$

which are the intersection of $\pi(\Sigma_{\pm})$. Taking account of $\pi(D)$ being one-dimensional, the boundary circles of D must be attached to C_a

$$\partial D = C_a \cup C_b. \quad (2.7)$$

In particular they satisfy

$$\pi(\partial D) = L_a \cup L_b. \quad (2.8)$$

On the other hand the vanishing of $\Omega_{\mathbf{R}}$ on D requires that D must lie in the hypersurface $b = \text{const.}$. Combining this requirement with eq.(2.7) we can conclude that all C_a must lie in this hypersurface for the vanishing of $\Omega_{\mathbf{R}}$ on D . Notice that this is not the condition on the membrane but the condition on the background geometry. It can be fulfilled by setting all the b -positions of the sixbranes equal. Let us specialize them to

$$b_1 = \dots = b_{N_f} = 0. \quad (2.9)$$

In this background geometry all C_a lie in the $b=0$ hypersurface of Q . One may suspect condition (2.9) loses a generality. But, from the viewpoint of the worldvolume gauge theory, the brane configuration under consideration describes the Coulomb branch and the b -direction is irrelevant in the description of this branch.

Now $\pi(D)$ is a one-dimensional line in the $b=0$ plane (which we will identify with the v -plane). It has the form

$$\pi(D) = L_a \cup L_{ab} \cup L_b \quad (2.10)$$

where L_{ab} is a one-dimensional line connecting L_a and L_b . Let us give somewhat an explicit description of L_a . Notice that the intersection of $\pi(\Sigma_{\pm})$ can be realized as the solution of the equation

$$|y_+(v)| = |y_-(v)|, \quad (2.11)$$

This equation tells that it is the region of the v -plane on which $\frac{\prod_a (v - \phi_a)^2}{\Lambda^{2N_c - N_f} \prod_j (v - e_j)}$ becomes a real non-negative quantity not bigger than 1. It consists of N_c one-dimensional lines connecting $v = \phi_a^{(+)}$ and $v = \phi_a^{(-)}$ through $v = \phi_a$. Here $\phi_a^{(\pm)}$ are the branch points of the curve which approach to ϕ_a as $\Lambda \rightarrow 0$. These one-dimensional lines are L_a .

Based on the form of $\pi(D)$ given in (2.10), we can say D has the following form in Q :

$$D = D_a \cup (L_{ab} \times S^1) \cup D_b, \quad (2.12)$$

where the S^1 is the circle of the eleven-dimension. D_a is the region of the cylinder $L_a \times S^1$ and is projected to L_a by π . The boundary of D_a consists of two circles one of which is identified with the boundary circle of the cylinder $L_{ab} \times S^1$ and the other is identified with the α -cycle C_a . Recall C_a is the circle wrapping once $L_a \times S^1$ and is projected to L_a by π . D_a occupies the half of $L_a \times S^1$ separated by C_a .

The area of the membrane is the sum of those of $D_{a,b}$ and $L_{ab} \times S^1$. The area of D_a is half the area of $L_a \times S^1$. Therefore it holds

$$A_D = A_{L_{ab} \times S^1} + \frac{1}{2}(A_{L_a \times S^1} + A_{L_b \times S^1}). \quad (2.13)$$

Let $L \times S^1$ be a cylinder in Q realized by $v=v(s), b=0$ and $\sigma=\sigma(t)$. s and t parametrize this cylinder. The induced metric h_{ij} of $L \times S^1$ becomes

$$\begin{aligned} h_{ss} &= \frac{R^2}{4} \left\{ V \left| \frac{dv}{ds} \right|^2 + \frac{1}{V} \left(\text{Im} \left(\delta \frac{dv}{ds} \right) \right)^2 \right\}, \\ h_{st} &= \frac{R^2}{4} \frac{1}{V} \text{Im} \left(\delta \frac{dv}{ds} \right) \frac{d\sigma}{dt}, \quad h_{tt} = \frac{R^4}{4} \frac{1}{V} \left(\frac{d\sigma}{dt} \right)^2. \end{aligned} \quad (2.14)$$

The volume form determined by this metric is

$$\begin{aligned} d\text{vol}_{L \times S^1} &= \frac{R^2}{4} \left| \frac{dv}{ds} \right| \frac{d\sigma}{dt} ds \wedge dt \\ &\equiv \frac{R^2}{4} |dv(s)| \wedge d\sigma(t). \end{aligned} \quad (2.15)$$

Therefore the area of $L \times S^1$ is $\pi R^2 |L|$. ($|L| \equiv \int_L |dv|$.) Now we can evaluate eq.(2.13) to

$$A_D = \pi R^2 \left(|L_{ab}| + \frac{|L_a| + |L_b|}{2} \right). \quad (2.16)$$

At this stage we want to minimize this A_D . Notice that L_a is fixed by the background geometry⁵. Only parameter which we can change is L_{ab} . It is the line connecting L_a and L_b . Clearly we must take the straight line for the minimization. The minimal area is

$$A_{min} = \pi R^2 \left(|\phi_a^{(-)} - \phi_b^{(+)}| + \frac{|L_a| + |L_b|}{2} \right). \quad (2.17)$$

But *this minimal area surface does not saturate the BPS bound*. This is because, though $\Omega_{\mathbf{R}}$ vanishes on D by the construction, the phase of Ω_C changes on $D_{a,b}$ and we can not tune it

⁵ M -theory brane configurations described in the previous section are treated as background geometry for membranes

to the definite phase which $\Omega_{\mathbf{C}}$ takes on $L_{ab} \times S^1$. The parts $D_{a,b}$ of the membrane are fixed by the background geometry. To achieve the BPS bound we need to change the background to a specific one by adjusting the values of the moduli parameters of the brane configuration. This will be discussed in the later subsection. But from the viewpoint describing the Coulomb branch, it is not favorable because of the loss of generality. Actually there exists another way to make the BPS saturated membrane. It is to consider a possibility of some recoil of the membrane to the background geometry. But to pursue such a possibility is beyond this paper.

The BPS bound has a nice interpretation [21, 22, 23] in terms of the Seiberg-Witten differential $\lambda_{SW} = v \frac{dy}{y}$ on the curve :

$$\left| \int_D \Omega_{\mathbf{C}} \right| = \frac{R^2}{2} \left| \oint_{C_a - C_b} \lambda_{SW} \right|. \quad (2.18)$$

Let us estimate the deviation of the minimal area from the BPS bound in the region $|\phi_a - \phi_b|, |\phi_a - e_j| \gg \Lambda$. One can estimate $|L_a|$ as $|L_a| \approx \Lambda$ in this region. Therefore the deviation is approximately

$$\frac{A_{min} - |\int_D \Omega_{\mathbf{C}}|}{|\int_D \Omega_{\mathbf{C}}|} \approx \frac{\Lambda}{|\phi_a - \phi_b|} \quad (2.19)$$

One can also discuss the mass of the minimal area membrane as an expansion by $\Lambda/|\phi_a - \phi_b|$. It has the form

$$\frac{A_{min}}{l_p^3} = \frac{\pi R^2}{l_p^3} |\phi_a - \phi_b| \left(1 + O \left(\frac{\Lambda}{|\phi_a - \phi_b|} \right) \right). \quad (2.20)$$

Rewriting the leading term by the original ten-dimensional coordinates it can be read as $\frac{\pi}{l_s^2} |(x_a^4 + ix_a^5) - (x_b^4 + ix_b^5)|$, where (x_a^4, x_a^5) are the positions of the fourbranes in the $x^{4,5}$ -directions. This is an expected result from Type IIA theory.

2.1.2 4-6 strings

We will consider a membrane which satisfies the second boundary condition. The background geometry is same as in the case of 4-4 strings. That is, $b_1 = \dots = b_{N_f} = 0$. Two-dimensional part D of the membrane is a cylinder which two boundary circles are attached respectively to the α -cycle C_a and the j -th sixbrane. For the vanishing of $\Omega_{\mathbf{R}}$ the cylinder D must lie in the hypersurface $b=0$. Following the notation of the previous study we can conclude D must have the following form in Q :

$$D = (L_{ja} \times S^1) \cup D_a \quad (\subset Q), \quad (2.21)$$

where L_{ja} is a one-dimensional line in the v -plane (which we identify with the $b=0$ plane of the $\mathbf{R}^3$) connecting $v=e_j$ and L_a . The area of D becomes

$$A_D = \pi R^2 \left(|L_{ja}| + \frac{|L_a|}{2} \right). \quad (2.22)$$

One can minimize the area by making L_{ja} the straight line. Again, notice that L_a is fixed by the background geometry. The minimal area is

$$A_{min} = \pi R^2 \left(|e_j - \phi_a^{(+)}| + \frac{|L_a|}{2} \right). \quad (2.23)$$

But, this minimal area surface does not saturate the BPS bound by the same reason as in the case of 4-4 strings. The BPS bound has a nice interpretation in terms of λ_{SW} besides the bare mass

$$\left| \int_D \Omega_{\mathbf{C}} \right| = \frac{R^2}{2} \left| 2\pi e_j - \oint_{C_a} \lambda_{SW} \right|. \quad (2.24)$$

The deviation of the minimal area from the BPS bound can be estimated in the region $|\phi_a - \phi_b|, |\phi_a - e_j| \gg \Lambda$

$$\frac{A_{min} - \left| \int_D \Omega_{\mathbf{C}} \right|}{\left| \int_D \Omega_{\mathbf{C}} \right|} \approx \frac{\Lambda}{|e_j - \phi_a|}. \quad (2.25)$$

In this region the mass of the minimal area membrane has the form

$$\frac{A_{min}}{l_p^3} = \frac{\pi R^2}{l_p^3} |e_j - \phi_a| \left(1 + O \left(\frac{\Lambda}{|e_j - \phi_a|} \right) \right). \quad (2.26)$$

Rewriting the leading part in terms of the original ten-dimensional coordinates it can be found to give an expected result from Type IIA theory.

2.1.3 6-6 strings

We will consider a membrane which satisfies the third boundary condition. The background geometry is same as in the previous cases. Two-dimensional part D of the membrane is a cylinder which two boundary circles are attached respectively to the j -th and k -th sixbranes. For the vanishing of $\Omega_{\mathbf{R}}$ the cylinder D must lie in the hypersurface $b=0$. Following the same notation as before D must have the following form in Q :

$$D = L_{jk} \times S^1 \quad (\subset Q), \quad (2.27)$$

where L_{jk} is a one-dimensional line in the v -plane connecting e_j and e_k . The area of D becomes

$$A_D = \pi R^2 |L_{jk}|. \quad (2.28)$$

One can minimize the area by making L_{jk} the straight line. The minimal area is

$$A_{min} = \pi R^2 |e_j - e_k|. \quad (2.29)$$

This minimal area surface saturate the BPS bound since $\Omega_{\mathbf{C}}$ has a constant phase on the straight line L_{jk} .

2.2 Dirichlet two-branes

In this case $\dim_{\mathbf{R}} D = 2$ holds. Let D be a two-dimensional disk. We can impose the following boundary condition on D

- The boundary circle of D is attached on Σ_{SW} in Q and, considering it as a one-dimensional cycle of Σ_{SW} , is homotopic to the β -cycle.

Membrane satisfying this boundary condition will gives rise to a D two-branes and provides a monopole in the worldvolume gauge theory.

This boundary condition is so severe in a general background geometry that we do not know how to minimize the area preserving it. The solution we have is very restrictive. Let us specialize [22] the background geometry to

$$\forall \phi_a, \quad \forall e_j \in \mathbf{R}. \quad (2.30)$$

Notice that the previous minimal area membranes corresponding to 4-4 and 4-6 strings also saturate their BPS bounds in this specific background geometry. This is simply because all L_a lie in the real line of the v -plane. In this background geometry one can take representatives of the β -cycles such that they are one-dimensional circles within the hypersurface $v = \bar{v}$ of Q . Let us denote them by B_a ($1 \leq a \leq N_c - 1$). One can easily see that σ takes constant values on these circles. Let D be a disk in the hypersurface $v = \bar{v}$ with its boundary being B_a and its σ -position being same as the boundary. Now on this disk $\Omega_{\mathbf{R}}$ vanishes and $\Omega_{\mathbf{C}}$ has a constant phase. So this membrane saturates the BPS bound as follows :

$$\begin{aligned} A_D &= \left| \int_D \Omega_{\mathbf{C}} \right| \\ &= \frac{R^2}{2} \left| \oint_{B_a} \lambda_{SW} \right|. \end{aligned} \quad (2.31)$$

3 Higgs Branch of $N=2$ MQCD

In this section we study the Higgs branch of $N=2$ MQCD. We first review the field theory results mainly based on [10]. Consider $N=2$ $SU(N_c)$ supersymmetric QCD with N_f flavors. We will treat the case of $N_f \leq N_c < 2N_f$. The classical vacua of this theory described by the F - and D -flat conditions. Let us denote the hypermultiplets by $(Q, \tilde{Q})$. $N=1$ chiral multiplets Q and $\tilde{Q}$ respectively belong to the fundamental and anti-fundamental representations of the gauge group $SU(N_c)$ besides the flavor group $U(N_f)$. We realize Q as a complex $N_c \times N_f$ matrix and $\tilde{Q}$ as a complex $N_f \times N_c$ matrix. In the Higgs branch the F - and D -flat conditions become

$$\begin{aligned} Q\tilde{Q} &= \zeta_{\mathbf{C}} \mathbf{1}_{N_c}, \\ QQ^\dagger - \tilde{Q}^\dagger \tilde{Q} &= \zeta_{\mathbf{R}} \mathbf{1}_{N_c}, \end{aligned} \quad (3.1)$$

where $\zeta_{\mathbf{C}, \mathbf{R}}$ are respectively complex and real numbers which we will call the Fayet-Iliopoulos (FI) parameters. $\vec{\zeta} \equiv (\zeta_{\mathbf{C}}, \zeta_{\mathbf{R}})$. Consideration on the Higgs branch is separated into two cases whether the FI parameters vanish or not. The former case is non-baryonic and the latter is baryonic. In the baryonic branch the color symmetry is completely broken due to the non-vanishing FI parameters. In the non-baryonic branch some of the color symmetry remains unbroken.

Let us examine the baryonic branch first. Fix the non-vanishing FI parameters at some values. Introducing the functions $\mu_{\mathbf{C}}(Q, \tilde{Q}) \equiv Q\tilde{Q}$ and $\mu_{\mathbf{R}}(Q, \tilde{Q}) \equiv QQ^\dagger - \tilde{Q}^\dagger \tilde{Q}$, we can write the moduli space of the (gauge equivalent) solutions of eqs.(3.1) as

$$\begin{aligned} \frac{\mu_{\mathbf{C}}^{-1}(\zeta_{\mathbf{C}}) \cap \mu_{\mathbf{R}}^{-1}(\zeta_{\mathbf{R}})}{SU(N_c)} &\simeq U(1) \times \frac{\mu_{\mathbf{C}}^{-1}(\zeta_{\mathbf{C}}) \cap \mu_{\mathbf{R}}^{-1}(\zeta_{\mathbf{R}})}{U(N_c)}, \\ &\equiv U(1) \times \hat{\mathcal{M}}_{\mathbf{b}}(\vec{\zeta}), \end{aligned} \quad (3.2)$$

where we rewrite the LHS using the isomorphism $U(N_c) \simeq U(1) \times SU(N_c)$. The second factor in this factorization, that is, the space $\hat{\mathcal{M}}_{\mathbf{b}}(\vec{\zeta})$ is a hyper-kähler manifold. With a little calculation one can find out it is a $4N_c(N_f - N_c)$ -dimensional smooth manifold. The FI parameters $\vec{\zeta}$ besides the $U(1)$ appearing in the above factorization can be also regarded as parameters of the baryonic branch. Let us denote their moduli space by P . Then the moduli space of the baryonic branch, which we denote by $\mathcal{M}_{\mathbf{b}}$, is the fibre bundle over P . The fibre space at $(\vec{\zeta}, e^{i\phi}) \in P$ is $\hat{\mathcal{M}}_{\mathbf{b}}(\vec{\zeta})$. So, $\mathcal{M}_{\mathbf{b}}$ is the moduli space of $N_c(N_f - N_c) + 1$ massless hypermultiplets.

Next, let us consider the non-baryonic branch. It is characterized by the vanishing FI parameters. The moduli space of the non-baryonic branch, which we denote by $\mathcal{M}_{\mathbf{nb}}$, have the

form :

$$\mathcal{M}_{\text{nb}} \equiv \frac{\mu_{\mathbf{C}}^{-1}(0) \cap \mu_{\mathbf{R}}^{-1}(0)}{U(N_c)}. \quad (3.3)$$

This is a singular manifold but it is possible to give a stratification by the color symmetry breaking patterns. For the breaking pattern, $SU(N_c) \rightarrow SU(N_c - r)$ we can introduce a stratum $\mathcal{M}_{\text{nb}}^{(r)}$ as the space of the (gauge equivalent) solutions of $\mu_{\mathbf{C}} = \mu_{\mathbf{R}} = 0$ which break only $U(r)$ color symmetry :

$$\mathcal{M}_{\text{nb}} = \cup_r \mathcal{M}_{\text{nb}}^{(r)}. \quad (3.4)$$

Each component $\mathcal{M}_{\text{nb}}^{(r)}$ can be considered as a hyper-Kähler manifold of dimensions $4r(N_f - r)$ and is called the moduli space of the r -th non-baryonic branch. So, this moduli space includes $r(N_f - r)$ massless hypermultiplets.

The Higgs branch intersects with the Coulomb branch. Intersection of the baryonic branch and the Coulomb branch is a single point of $\mathcal{M}_{\text{coul}}$, the moduli space of the Coulomb branch. This is called the baryonic branch root. Though $U(1)^{N_c-1}$ color symmetry remains unbroken at a generic point of the Coulomb branch, it must be completely broken at this root. This is due to the requirement from the baryonic branch side. The intersection of the r -th non-baryonic branch and the Coulomb branch is a $2(N_c - r - 1)$ -dimensional submanifold of $\mathcal{M}_{\text{coul}}$. This is called the r -th non-baryonic branch root. On this submanifold only $U(1)^{N_c-r-1}$ color symmetry remains unbroken. This is because $U(r)$ color symmetry must be broken by the requirement from the r -th non-baryonic branch side and the remaining $SU(N_c - r)$ color symmetry is broken down to $U(1)^{N_c-r-1}$ of the low energy effective theory.

In the following subsections we present the descriptions of the baryonic and non-baryonic branches in terms of the M -theory brane configurations. We follow the argument given in [17]. We begin by studying their roots using the M -theory description of the Coulomb branch in the previous sections. In that description, the Seiberg-Witten curve, that is, two-dimensional part of the M fivebrane, is holomorphically embedded into the multi-Taub-NUT space Q , which is the geometry sixbranes provide in M -theory. Moduli parameters of this configuration are $\phi_a \in \mathbf{C}$ for the fivebrane and $(e_i, b_i) \in \mathbf{C} \times \mathbf{R}$ for the sixbranes (or the multi-Taub-NUT space). After specifying these moduli parameters at the values relevant for the roots, we will show that the baryonic and non-baryonic branches are actually realized in M -theory.

3.1 Non-baryonic Branch

We consider the r -th non-baryonic branch root, that is the intersection of $\mathcal{M}_{\text{nb}}^{(r)}$ and $\mathcal{M}_{\text{coul}}$, from the Coulomb branch side. The corresponding brane configuration is given by the conditions,

$$\phi_1 = \cdots = \phi_r = 0, \quad \phi_a \neq 0 \quad (r+1 \leq a \leq N_c) \quad (3.5)$$

for the fivebrane and

$$e_1 = \cdots = e_{N_f} = 0 \quad (3.6)$$

for the sixbranes. We set the b -positions of the sixbranes as follows :

$$b_1 < \cdots < b_{N_f}. \quad (3.7)$$

If one takes Type IIA picture it will correspond to the configuration in which r of N_c D fourbranes overlap and N_f D sixbranes cross these r D fourbranes. Their crossing breaks the $U(r)$ color symmetry completely.

To study this configuration let us recall compact two-cycles of Q . In general there are $N_f - 1$ topologically nontrivial two-spheres in Q . Let us denote them by S_j . By the natural projection $\pi: Q \rightarrow \mathbf{R}^3$, each two-sphere S_j is projected to the one-dimensional straight line l_j connecting (e_j, b_j) and (e_{j+1}, b_{j+1}) . The symplectic two-forms $\Omega_{\mathbf{R}, \mathbf{C}}$ take the following form on S_j :

$$\Omega_{\mathbf{R}}|_{S_j} = \frac{R^2}{4} db \wedge d\sigma, \quad \Omega_{\mathbf{C}}|_{S_j} = \frac{R^2}{4} dv \wedge d\sigma. \quad (3.8)$$

In particular, at the configuration relevant for the non-baryonic branch root, the holomorphic two-form restricted on S_j vanishes. This means that all S_j are holomorphic in Q at the root.

At the non-baryonic branch root one can expect that Σ_{SW} wraps these two-cycles S_j . Wrapping and unwrapping parts of Σ_{SW} intersect only at the points where the sixbranes are located. We can regard these wrapping parts of the fivebrane as fivebranes with worldvolume $\mathbf{R}^4 \times S_j$. They can move to the $x^{7,8,9}$ -directions on the sixbranes without any obstruction. Rotation group of these directions is identified with $SU(2)_R$ of $N = 2$ supersymmetry. So, they will provide [2] massless hypermultiplets in the non-baryonic branch.

How many massless hypermultiplets can we obtain? To count their number and to check that Σ_{SW} actually wraps these two-cycles, it is convenient to study the holomorphic embedding of the curve in a different fashion,

$$y(v, b, \sigma) + z(v, b, \sigma) = 2v^r \prod_{a=r+1}^{N_c} (v - \phi_a), \quad (3.9)$$

where $z(v, b, \sigma) \equiv \Lambda^{2N_c - N_f} v^{N_f} / y(v, b, \sigma)$.

To see whether Σ_{SW} wraps S_j or not, it is enough to show the following : For any b^* satisfying $b_j < b^* < b_{j+1}$, $v=0$ is a solution of the embedding equation at b^* ,

$$y(v, b^*, \sigma) + z(v, b^*, \sigma) = 2v^r \prod_{a=r+1}^{N_c} (v - \phi_a). \quad (3.10)$$

Let us explain why this is enough. Suppose $v=0$ is a solution of eq.(3.10) for any b^* . Then $l_j (= \pi(S_j))$ is included in $\pi(\Sigma_{SW})$. From the dimensionality this means that S_j is included in Σ_{SW} . Taking this viewpoint the multiplicity of solution $v=0$ for eq.(3.10) is nothing but the wrapping number of Σ_{SW} to S_j . This is the number of the fivebranes with worldvolume $\mathbf{R}^4 \times S_j$.

Let us show the above claim. Consider eq.(3.10) in the region $|v| \ll |b^* - b_j|, |b^* - b_{j+1}|$. In this region $y(v, b^*, \sigma)$ and $z(v, b^*, \sigma)$ behave as

$$\begin{aligned} y(v, b^*, \sigma) &\sim \prod_{i=1}^{N_f} (-b^* + b_i + |b^* - b_i| + \frac{1}{2}|v|^2/|b^* - b_i|)^{1/2}, \\ z(v, b^*, \sigma) &\sim \prod_{i=1}^{N_f} (b^* - b_i + |b^* - b_i| + \frac{1}{2}|v|^2/|b^* - b_i|)^{1/2}. \end{aligned} \quad (3.11)$$

Notice that the following equalities

$$\begin{aligned} -b + b_i + |b - b_i| &= \begin{cases} 2(-b + b_i) & (\neq 0) \text{ if } b < b_i \\ 0 & \text{if } b > b_i \end{cases}, \\ b - b_i + |b - b_i| &= \begin{cases} 0 & \text{if } b < b_i \\ 2(b - b_i) & (\neq 0) \text{ if } b > b_i \end{cases}. \end{aligned}$$

So, eqs.(3.11) imply

$$\begin{aligned} y(v, b^*, \sigma) &\sim v^j \times (\text{non-vanishing factor at } v=0), \\ z(v, b^*, \sigma) &\sim v^{N_f - j} \times (\text{non-vanishing factor at } v=0). \end{aligned} \quad (3.12)$$

Now eq.(3.10) acquires the following form in this region

$$v^j + v^{N_f - j} = v^r. \quad (3.13)$$

This means that $v=0$ is a solution of eq.(3.10) for $b_j < b^* < b_{j+1}$. Its multiplicity, which we denote by n_j , can be read from eq.(3.13)

$$n_j = \min \{j, N_f - j, r\}. \quad (3.14)$$

This coincides with the result derived in Type IIA picture using the s -rule [7]. Actually the above argument gives a proof of the s -rule via M-theory. Since all S_j are holomorphic at the root one can also show [24] the s -rule as the resolution of the curve. Now the total number of the massless hypermultiplets is the sum of n_j

$$\sum_{j=1}^{N_f-1} n_j = r(N_f - r). \quad (3.15)$$

This is the correct number of massless hypermultiplets allowed in the r -th non-baryonic branch.

3.2 Baryonic Branch

The baryonic branch root, that is, the intersection of $\mathcal{M}_b$ and $\mathcal{M}_{\text{coul}}$, is a single point of $\mathcal{M}_{\text{coul}}$ where the color symmetry $SU(N_c)$ is completely broken. At a first glance one may expect it is given by $\forall \phi_a = \forall e_j = 0$. But this naive expectation is incorrect in quantum theory. Due to the quantum correction on the Coulomb branch the root changes from the naive one. Where is the root? We first claim that the curve is completely degenerate at the root. This is because $SU(N_c)$ must be broken at the root. In quantum theory $U(1)_R$ symmetry is broken to $\mathbf{Z}_{2(2N_c-N_f)}$ by the chiral anomaly, which acts on the v -plane as $\mathbf{Z}_{2N_c-N_f}$. So, we also claim that the curve at the root is invariant under this discrete symmetry. With these two claims the baryonic branch root was determined [10]. It is given by

$$\begin{aligned} \phi_1 = \cdots = \phi_{N_f-N_c} = 0, \quad \phi_r = \omega^{r-N_f+N_c} \Lambda \quad (N_f - N_c + 1 \leq r \leq N_c), \\ e_1 = \cdots = e_{N_f} = 0, \end{aligned} \quad (3.16)$$

where $\omega \equiv e^{\frac{2\pi i}{2N_f-N_c}}$. The curve at the root becomes

$$y^2 - 2v^{N_f-N_c}(v^{2N_c-N_f} - \Lambda^{2N_c-N_f})y - 4\Lambda^{2N_c-N_f}v^{N_f} = 0. \quad (3.17)$$

It is invariant under the $\mathbf{Z}_{2N_c-N_f}$ symmetry and is completely factorized :

$$y_+(v) = 2v^{N_c}, \quad y_-(v) = -2\Lambda^{2N_c-N_f}v^{N_f-N_c}. \quad (3.18)$$

The previous consideration on the wrapping of Σ_{SW} to the two-spheres S_j works also on this case. Setting r in (3.14) to be $N_f - N_c$, the wrapping number n_j of Σ_{SW} to S_j is

$$n_j = \min \{j, N_f - j, N_f - N_c\}. \quad (3.19)$$

So, the total number of the massless hypermultiplets given by the fivebranes with worldvolume $\mathbf{R}^4 \times S_j$ is

$$\sum_{j=1}^{N_f-1} n_j = N_c(N_f - N_c). \quad (3.20)$$

This is the correct number of massless hypermultiplets in $\hat{\mathcal{M}}_{\mathbf{b}}(\vec{\xi})$.

4 Duality of Baryonic Branch

The baryonic branch root is located in the strong coupling region of the original microscopic $SU(N_c)$ gauge theory. Gauge theory description in the neighborhood of the root can be obtained by applying electric-magnetic duality. The magnetic theory is [10] the $SU(N_f - N_c) \times U(1)^{2N_c - N_f}$ gauge theory which has $2N_c - N_f$ massless singlet hypermultiplets charged only by the $U(1)$ factors in addition to N_f massless quark hypermultiplets belonging to the fundamental representations of $SU(N_f - N_c)$. Baryonic branch of this IR-effective theory exactly coincides [10] with that of the original microscopic $SU(N_c)$ theory. We will call the original microscopic and IR-effective theories respectively “electric” and “magnetic” ones.

Let us first consider Type IIA brane configuration relevant to the magnetic theory. We need N_c D fourbranes (suspended by two NS fivebranes) to obtain the vector multiplets. We also need $2N_c$ D sixbranes to obtain the hypermultiplets. Notice that the $2N_c - N_f$ singlet hypermultiplets and the N_f quark hypermultiplets in the magnetic theory are elementary. These hypermultiplets should appear in the light spectrum of 4-6 strings. To obtain the non-abelian part of the magnetic theory, $N_f - N_c$ fourbranes must be overlapping and N_f sixbranes must be crossing these overlapping fourbranes. To obtain the abelian part, $2N_c - N_f$ fourbranes must be separated from each other. Each one of them must be crossed by a single sixbrane. The positions of the separated fourbranes must be invariant under the $\mathbf{Z}_{2N_c - N_f}$ discrete symmetry. This is necessary to provide the magnetic description of the baryonic branch root.

At this stage we encounter some puzzle. Type IIA brane configuration relevant to the electric description of the baryonic branch root consists of N_c fourbranes and N_f sixbranes. All the fourbranes must be overlapping and all the sixbranes must be crossing these fourbranes. Notice that this configuration is allowed by the s -rule. Electric and magnetic brane configurations are different from each other although they describe the same root. The difference originates in the existence of the extra $2N_c - N_f$ sixbranes of the magnetic configuration. In the electric




Figure 2: Creation and annihilation of D fourbrane by D sixbrane in Type IIA picture.

description the corresponding singlet hypermultiplets are considered as monopoles. So, they are not elementary.

Let us solve this puzzle in Type IIA theory. It is argued in [7] that, when two NS fivebranes are exchanged crossing a D sixbrane, a D fourbrane is created and then suspended between these fivebranes with touching the sixbrane. Conversely, a D fourbrane suspended between these fivebranes with touching the sixbrane is annihilated by this exchange. See Fig.2. With this process of the exchange the extra $2N_c - N_f$ sixbranes in the magnetic configuration annihilate the $2N_c - N_f$ fourbranes touching them. The N_f sixbranes which are crossing the overlapping $N_f - N_c$ fourbranes create N_c fourbranes to preserve the s -rule. See Fig.3. So, after the exchange, the magnetic configuration becomes very similar to the electric configuration. Finally, letting the extra $2N_c - N_f$ sixbranes go to infinity, we actually obtain the electric configuration.

We want to make the above Type IIA explanation on the duality between the electric and magnetic descriptions of the baryonic branch root more rigorous by taking the M -theory viewpoint. In the next subsection we will explain the brane creation by considering a M -theory configuration consisting of a fivebrane and a sixbrane. This explanation is due to [25]. Roughly speaking, different boundaries of the moduli space of the M -theory configuration will give rise to different IIA configurations. They will be distinguished by the appearance of a D fourbrane. In the last subsection, following [26], we present a family of M -theory configurations relevant to describe the brane exchange associated with the above duality.

4.1 Brane creation via M -theory

Consider a M -theory configuration consisting of a single fivebrane and a single sixbrane. Two-dimensional part of the fivebrane is holomorphically embedded to the Taub-NUT space which the sixbrane provides. Let $(v, b) = (0, -b_0)$ be the position of the sixbrane. Holomorphic




Figure 3: Brane configurations related by the exchange of the NS fivebranes. (a) is “electric” and (b) is “magnetic”. The case of $(N_c, N_f) = (4, 6)$ is sketched as an example.

embedding of the fivebrane can be described by

$$y(v, b, \sigma) = v, \quad (4.1)$$

where $y(v, b, \sigma)$ is the holomorphic coordinate of the Taub-NUT space

$$y(v, b, \sigma) = e^{-\frac{b+i\sigma}{2}} \left(-b - b_0 + \sqrt{(b + b_0)^2 + |v|^2} \right). \quad (4.2)$$

We examine the above embedding equation by separating it into its angular and radial parts. The angular part becomes

$$e^{-\frac{\sigma}{2}} = \frac{v}{|v|}, \quad (4.3)$$

which describes a vortex of σ at $v=0$. Recall that, in order to introduce the complex structure, we fix the Dirac string of the sixbrane such that it runs from $(0, -b_0)$, parallel with the b -axis, to $+\infty$. The vortex of σ describes the intersection of the fivebrane and the Dirac string. It means that *the fivebrane is sitting to the right of the sixbrane in the $\mathbf{R}^3$ (or the Taub-NUT space)*. Fivebrane sitting to the left of the sixbrane can be described by

$$z(v, b, \sigma) = v, \quad (4.4)$$

where $z(v, b, \sigma) \equiv v/y(v, b, \sigma)$ ⁶.

The radial part of eq.(4.1), after a parallel shift of the b -coordinate, acquires the form

$$e^{-b} = e^{-b_0} \left(b + \sqrt{b^2 + |v|^2} \right). \quad (4.5)$$

⁶An interesting observation on the asymmetry between eqs.(4.1) and (4.4) was made in [27].




Figure 4: Shapes of the fivebrane in the $(|v|, b)$ -plane.

Regarding it as an equation for b , the solution does depend only on $|v|$ and b_0 . Denote the solution by $b(|v|; b_0)$. Its independence from the phase of v means that the fivebrane has a rotational symmetry around the b -axis in the $\mathbf{R}^3$. Differentiating eq.(4.5) by $|v|$ one can easily find that $b(|v|; b_0)$ is monotonically decreasing w.r.t $|v|$. The shapes of the curve $b = b(|v|; b_0)$, somewhat dependent on the value of b_0 , are depicted in Fig.4

There exists three characteristic points on this curve. The first one is the intersection with the b -axis. The second is the intersection with the line $b=0$. The last is the intersection with the line $b = b_0$, which exists only for the case $b_0 \leq 1/2$. These three points on the curve are denoted by A, B and C. (See Fig.4.) Their coordinates in the $(|v|, b)$ -plane can be read from eq.(4.5)

$$A : (|v|, b) = (0, \beta(b_0)), \quad B : (e^{b_0}, 0), \quad C : (\sqrt{1 - 2b_0}, b_0), \quad (4.6)$$

where we introduce the function $\beta(b_0)$ by $2\beta e^\beta = e^{b_0}$.

We want to consider the IIA limit of this M -theory brane configuration. It will be achieved [28] by taking $R \rightarrow 0$ with fixing the value of $Rb_0/2$. Let us denote this fixed value by x_0^6 . Recall that, back to the original ten-dimensional coordinates, the value of $Rb_0/2$ provides the x^6 -position of the sixbrane. (The $x^{4,5}$ -positions have been set to zero from the beginning.) So, this limit simply means making the radius of the eleven-dimension circle very small fixing the position of sixbrane. In this course of procedure, the fivebrane is always holomorphically embedded into the Taub-NUT space preserving its BPS saturation. So, in the limit we can expect to obtain a BPS saturated object of Type IIA theory.

In order to fix $x_0^6 = Rb_0/2$ at a positive value for small R , b_0 must be a large positive quantity. While, in order to fix x_0^6 at a negative value for small R , b_0 must be a large negative quantity. Let us first examine the case of $x_0^6 < 0$. Consider the fivebrane in the $(|v|, b)$ -plane. It is given by the curve $b = b(|v|; b_0)$. Back to the original ten-dimensional coordinates, its coordinates become $(|x^4 + ix^5|, x^6) = \frac{R}{2}(|v|, b)$. We want to know how the curve in the original coordinates behaves under the limit. Let us pay attention to the characteristic points A, B and C on the curve. Quantities appearing in (4.6) turn out to become as follow : $\lim'_{R \rightarrow 0} R\beta(b_0) = \lim'_{R \rightarrow 0} Re^{b_0} = \lim'_{R \rightarrow 0} R\sqrt{1 - 2b_0} = 0$, where “ $\lim'_{R \rightarrow 0}$ ” means taking $R \rightarrow 0$ with fixing $x_0^6 = Rb_0/2$. So, A, B and C become the following points on the $(|x^4 + ix^5|, x^6)$ -plane

$$A : (|x^4 + ix^5|, x^6) = (0, 0), \quad B : (0, 0), \quad C : (0, x_0^6). \quad (4.7)$$

While this, one can also see that $\partial b(|v|; b_0)/\partial |v|$ goes to zero under this limit. Gathering all these information, we can conclude that the segment of the curve between A and C (which includes B) approaches to the straight line between $(0, x_0^6)$ and $(0, 0)$ and that the residual part of the curve goes to the line $x^6 = x_0^6$. See Fig.5 (a). Owing to the rotational symmetry of the curve around the x^6 -axis the line $x^6 = x_0^6$ realizes a five-dimensional object in ten-dimensions, that is, a NS fivebrane while the line segment between $(0, x_0^6)$ and $(0, 0)$ realizes a four-dimensional object in ten-dimensions, that is, a fourbrane. To summary, the IIA limit of the M -theory configuration with b_0 negative describes the configuration of a NS fivebrane, a D fourbrane and a D sixbrane. The NS fivebrane is located at $(x^6, x^7, x^8, x^9) = (x_0^6, 0, 0, 0)$ while the D fourbrane with worldvolume $(x^0, x^1, x^2, x^3, x^6)$ is suspended, in the (x^4, x^5, x^6) -space, between the point $(0, 0, x_0^6)$ on the NS fivebrane and the D sixbrane at $(0, 0, 0)$.

Next let us consider the case of $x_0^6 > 0$. We can repeat the same argument. Consider the characteristic points A and B on the curve. Notice that, since we can assume $b_0! > 1/2$ without losing any generality, the point C is irrelevant for the discussion. Quantities appearing in (4.6) have the following limits in this case : $\lim'_{R \rightarrow 0} R\beta(b_0) = 2x_0^6$ and $\lim'_{R \rightarrow 0} Re^{b_0} = +\infty$. So, A and B become the following points on the $(|x^4 + ix^5|, x^6)$ -plane

$$A : (|x^4 + ix^5|, x^6) = (0, x_0^6), \quad B : (+\infty, 0). \quad (4.8)$$

Since it holds also $\partial b(|v|; b_0)/\partial |v| \rightarrow 0$, we can conclude that the curve approaches to the straight line $x^6 = x_0^6$. So, we obtain only a fivebrane. See Fig.5 (b). To summarize, the IIA limit of the M -theory configuration with b_0 positive describes the configuration of a NS fivebrane and a D sixbrane. The NS fivebrane is located at $(x^6, x^7, x^8, x^9) = (x_0^6, 0, 0, 0)$ and the D sixbrane is at $(x^4, x^5, x^6) = (0, 0, 0)$. There appears no D fourbrane.




Figure 5: IIA limits of M -theory brane configuration depicted in Fig.4: (a) D fourbrane is suspended between NS fivebrane and D sixbrane. (b) There are only NS fivebrane and D sixbrane.

For the M -theory configuration (4.1) with the moduli parameter b_0 we have shown that there are two possibilities of the IIA configuration, one of which is the configuration of a NS fivebrane, a D fourbrane and a D sixbrane while the other is the configuration of a NS fivebrane and a D sixbrane. Appearance or disappearance of the D fourbrane in the IIA configuration can be understood as a result of the crossing of NS fivebrane and D sixbrane if one takes Type IIA picture.

4.2 Duality of baryonic branch via M-theory

Let us consider the following family of $\mathbf{Z}_{2N_c-N_f}$ -invariant curves Σ_ρ

$$y^2 - 2v^{N_f-N_c}(v^{2N_c-N_f} - \Lambda^{2N_c-N_f})y + f(\rho)v^{N_f}(v^{2N_c-N_f} - g(\rho)\Lambda^{2N_c-N_f}) = 0. \quad (4.9)$$

Here $\rho \in [0, 1]$ parametrizes the family. f and g are some unknown functions of ρ . To determine them we require the following constraints on the curves :

- As $\rho \rightarrow 0$, Σ_ρ becomes the Seiberg-Witten curve for the electric description of the baryonic branch root while, as $\rho \rightarrow 1$, Σ_ρ becomes the Seiberg-Witten curve for the magnetic description of the baryonic branch root.
- Σ_ρ is completely degenerate for any ρ .

These requirements are based on a naive M-theory generalization of the exchange of the NS fivebranes in Type IIA theory. The first one is, so to say, the initial and final conditions of

the brane exchange. And the second one reflects the fact that the NS fivebranes must be disconnected under their exchange.

These constraints put some restrictions on f and g . The complete degeneration of Σ_ρ requires f be the following function of g

$$f = \frac{4(g-1)}{g^2}. \quad (4.10)$$

The initial and final conditions restrict the boundary behaviors of g as

$$\lim_{\rho \rightarrow 0} g(\rho) = \infty, \quad \lim_{\rho \rightarrow 1} g(\rho) = 1. \quad (4.11)$$

In principle we can take any function g with these boundary behaviors. But, this arbitrariness is irrelevant. It simply means that one can move the fivebranes in an arbitrary manner as far as they exchange each other. We will fix it as

$$g(\rho) = \frac{1}{\rho}. \quad (4.12)$$

With this choice of g the factorized form of the curve Σ_ρ becomes

$$y_+(v) = 2(1-\rho)v^{N_c}, \quad y_-(v) = 2\rho v^{N_f-N_c}(v^{2N_c-N_f} - \Lambda^{2N_c-N_f}/\rho). \quad (4.13)$$

These curves are holomorphically embedded into the multi Taub-NUT spaces. The multi Taub-NUT space Q_ρ in which Σ_ρ is embedded is characterized by the positions of $2N_c$ sixbranes, (e_i, b_i) . Their v -positions are specified from the equation of the curve

$$e_1 = \cdots = e_{N_f} = 0, \quad e_{N_f+r}(\rho) = \omega^r \Lambda / \rho^{\frac{1}{2N_c-N_f}} \quad (1 \leq r \leq 2N_c - N_f). \quad (4.14)$$

We will fix their b -positions to $b_1 = \cdots = b_{2N_c} = 0$.

The holomorphic embedding of Σ_ρ into Q_ρ is given by

$$y_\pm(v) = y(v, b, \sigma). \quad (4.15)$$

Since the curve Σ_ρ is completely factorized, each equation describes independently the embedding of a complex plane $\Sigma_{\rho+}$ or $\Sigma_{\rho-}$. Consider the asymptotic b -positions of $\Sigma_{\rho\pm}$, that is, the values of $b_\pm(v)$ at $v \approx \infty$. A simple estimation of eqs.(4.15) gives $b_+(v) \approx -2 \ln 2(1-\rho) - 2N_c \ln |v|$ and $b_-(v) \approx -2 \ln 2\rho - 2N_c \ln |v|$. Therefore their relative b -position at $v = \infty$ is $b_+(\infty) - b_-(\infty) = 2 \ln \frac{\rho}{1-\rho}$, which is negative when $0 < \rho < 1/2$ and positive when $1/2 < \rho < 1$. This means that $\Sigma_{\rho\pm}$ overlap asymptotically at $\rho = 1/2$. We want to propose that the brane exchange actually occurs at $\rho = 1/2$.

To justify this proposal we discuss massless membranes associated with the fivebrane at each value of ρ . Let us begin by studying the intersection of $\Sigma_{\rho\pm}$ in the $\mathbf{R}^3$. Let π be the natural projection from Q_ρ to $\mathbf{R}^3$ as before. The intersection $\pi(\Sigma_{\rho+}) \cap \pi(\Sigma_{\rho-})$ is located at the $b=0$ plane in $\mathbf{R}^3$, which we identify with the v -plane. The intersection is the solution of the equation

$$|y_+(v)| = |y_-(v)|. \quad (4.16)$$

Using the explicit forms (4.13) it is solved as

$$v_r(\theta) = \frac{e_{N_f+r}(\rho)}{\left(1 + e^{i\theta} \left| \frac{1-\rho}{\rho} \right| \right)^{\frac{1}{2N_c-N_f}}}, \quad (r = 1, \dots, 2N_c - N_f, \quad 0 \leq \theta \leq 2\pi.) \quad (4.17)$$

So, the intersection consists of one-dimensional lines l_r given by $v=v_r(\theta)$

Actual forms of the lines l_r drastically change at $\rho=1/2$. When $0 < \rho < 1/2$, each line l_r is connected with $l_{r\pm 1}$ at the boundaries. $l_1 \cup \dots \cup l_{2N_c-N_f}$ becomes a single closed loop encircling $v=0$. On the other hand, when $1/2 < \rho < 1$, each line l_r itself becomes a closed loop on the v -plane encircling $v=e_{N_f+r}(\rho)$.

What does this qualitative difference of the intersection mean? To explain this we shall consider a slight perturbation of Σ_ρ . We slightly change the v -positions of the $2N_c-N_f$ sixbranes from $v=e_{N_f+r}(\rho)$. With this change the curve Σ_ρ becomes $\tilde{\Sigma}_\rho$. This perturbed curve has genus $2N_c-N_f$. Its Riemann sheets $\tilde{\Sigma}_{\rho\pm}$ are patched together by $2N_c-N_f$ circles $\tilde{C}_r$ in the multi-Taub-NUT space. Each $\tilde{L}_r \equiv \pi(\tilde{C}_r)$ is a one-dimensional line in the v -plane connecting two branch points of $\tilde{\Sigma}_\rho$ through $v=\omega^r \Lambda$.

We first examine the case of $0 < \rho < 1/2$. The ends of $\tilde{L}_r$ (two branch points of $\tilde{\Sigma}_\rho$) are getting closer to the ends of $\tilde{L}_{r\pm 1}$ as the perturbation becomes weak. Finally they are identified with the ends of $\tilde{L}_{r\pm 1}$. It means that $2N_c-N_f$ one-dimensional lines $\tilde{L}_r$ become the single closed loop $l_1 \cup \dots \cup l_{2N_c-N_f}$. In this process the β -cycles of $\tilde{\Sigma}_\rho$ vanish. Nextly we consider the case of $1/2 < \rho < 1$. The ends of $\tilde{L}_r$ are getting closer to each other as the perturbation becomes weak. Finally its ends are identified with each other. So, we obtain the closed loop l_r . In this process the α -cycles of $\tilde{\Sigma}_\rho$ vanish.

The above consideration implies that the complete degeneration of Σ_ρ is due to the vanishing $2N_c-N_f$ β -cycles when $0 < \rho < 1/2$ and due to the vanishing $2N_c-N_f$ α -cycles when $1/2 < \rho < 1$. This reflects the qualitative difference of the intersection in the $\mathbf{R}^3$.

Now we consider massless membranes. First we examine the case of $1/2 < \rho < 1$. Consider a membrane with worldvolume $\mathbf{R} \times D$, where D is a disk and its boundary is on Σ_ρ such that

$\pi(D)$ is in the v -plane encircled by l_r . This membrane also touches the sixbrane at $v = e_{N_f+r}(\rho)$. We can deform this membrane keeping its boundary on Σ_ρ and touching the sixbrane. Actually we can make the area arbitrary small. This is because the boundary circle of D can shrink to a point of Σ_ρ due to the vanishing of the α -cycles. So the minimal area is zero. It is achieved not by a membrane but by a string. This string provides a massless hypermultiplet. It is a singlet hypermultiplet in the magnetic description of the baryonic branch root. Next we examine the case of $0 < \rho < 1/2$. Consider the curve $\tilde{\Sigma}_\rho$ and imagine a membrane with worldvolume $\mathbf{R} \times D$ where D is a disk with its boundary is one of the β -cycles of $\tilde{\Sigma}_\rho$. Since Σ_ρ is obtained by vanishing the β -cycles of $\tilde{\Sigma}_\rho$, the area of this membrane becomes zero after the degeneration. In this limit D can be regarded as a point. This provides a massless monopole in the electric description of the baryonic branch.

References

- [1] A. Giveon and D. Kutasov, "Brane Dynamics and Gauge Theory", hep-th/9802067 and references there in.
- [2] E. Witten, "Solutions of Four-Dimensional Field Theories via M Theory", hep-th/9703166.
- [3] N. Seiberg and E. Witten, "Electric-Magnetic Duality, Monopole Condensation, and Confinement in N=2 Supersymmetric Yang-Mills Theory," Nucl. Phys. **B426** (1994) 19, hep-th/9407087.
- [4] N. Seiberg and E. Witten, "Monopoles, Duality and Chiral Symmetry Breaking in N=2 Supersymmetric QCD," Nucl. Phys. **B431** (1994) 484, hep-th/9408099.
- [5] N. Seiberg, "Electric-Magnetic Duality in Supersymmetric Non-Abelian Gauge Theories", Nucl. Phys. **B435** (1995) 129, hep-th/9411149.
- [6] K. Intriligator and N. Seiberg, "Lectures on Supersymmetric Gauge Theories and Electric-Magnetic Duality", Nucl. Phys. Proc. Suppl. **45BC** (1996) 1, hep-th/9509066.
- [7] A. Hanany and E. Witten, "Type IIB Superstrings, BPS Monopoles, and Three-dimensional Gauge Dynamics", Nucl. Phys. **B492** (1997) 152, hep-th/9611230.
- [8] S. Elitzur, A. Giveon and D. Kutasov, "Branes and N=1 Duality in String Theory", Phys. Lett. **400B** (1997) 269, hep-th/9702014.

- [9] E. Witten, "String Theory Dynamics in Various Dimensions", Nucl. Phys. **B443** (1995) 85, hep-th/9503124.
- [10] P. C. Argyres, M. R. Plesser and N. Seiberg, "The Moduli Space of Vacua of $N=2$ SUSY QCD and Duality in $N=1$ SUSY QCD," Nucl. Phys. **B471** (1996) 159, hep-th/9603042.
- [11] E. Verlinde, "Global Aspects of Electric - Magnetic Duality", Nucl. Phys. **B455** (1995) 211, hep-th/9506011.
- [12] A. Klemm, W. Lerche, S. Yankielowicz, "Simple Singularities and $N = 2$ Supersymmetric Yang-Mills Theory," Phys. Lett. **B344** (1995) 169, hep-th/9411048.
- [13] P. C. Argyres and A. E. Faraggi, "The Vacuum Structure and Spectrum of $N = 2$ Supersymmetric $SU(N)$ Gauge Theory," Phys. Rev. Lett. **74** (1995) 3931, hep-th/9411057.
- [14] G.T. Horowitz and A. Strominger, "Black strings and p-Branes", Nucl. Phys. **B360** (1991) 197
- [15] P. K. Townsend, "The eleven-dimensional supermembrane revisited", Phys. Lett. **B350** (1995) 184, hep-th/9501068.
- [16] S. W. Hawking, "Gravitational Instantons," Phys. Lett. **60A** (1977) 81.
G. W. Gibbons and S. W. Hawking, "Classification of Gravitational Instanton Symmetries", Commun. Math. Phys. **66** (1979) 291.
- [17] T. Nakatsu, K. Ohta, T. Yokono and Y. Yoshida, "Higgs Branch of $N=2$ SQCD and M Theory Branes", hep-th/9707258.
- [18] N. J. Hitchin, "Polygons and gravitons", Math. Proc. Camb. Phil. Soc. **85** (1979) 465.
- [19] A. Hanany and Y. Oz, "On the Quantum Moduli Space of Vacua of $N = 2$ Supersymmetric $SU(N_c)$ Gauge Theories," Nucl. Phys. **B452** (1995) 283, hep-th/9505075.
- [20] P. C. Argyres, M. R. Plesser and A. D. Shapere, "The Coulomb Phase of $N=2$ Supersymmetric QCD," Phys. Rev. Lett. **75** (1995) 1699, hep-th/9505100.
- [21] A. Fayyazuddin and M. Spalinski, "The Seiberg-Witten Differential from M Theory", hep-th/9706087.
- [22] M. Henningson and P. Yi, "Four-dimensional BPS-spectra via M-theory", hep-th/9707251.

- [23] A. Mikhailov, “BPS States and Minimal Surfaces”, hep-th/9708068.
- [24] K. Hori, H. Ooguri and Y.Oz, “Strong Coupling Dynamics of Four-Dimensional N=1 Gauge Theories from M theory Fivebrane”, hep-th/9706082.
- [25] T. Nakatsu, K. Ohta, T. Yokono and Y. Yoshida, “A Proof of Brane Creation via M theory”, hep-th/9711117.
- [26] T. Nakatsu, K. Ohta and T. Yokono, ”On the Baryonic Branch Root of N=2 MQCD”, hep-th/9712029.
- [27] C.P. Bachas and M.B. Green, ”A Classical Manifestation of the Pauli Exclusion Principle”, hep-th/9712187.
- [28] A. Brandhuber, N. Itzhaki, V. Kaplunovsky, J. Sonnenschein and B. S. Yankielowicz, ”Comments on the M theory Approach to N=1 SQCD and Brane Dynamics”, hep-th/9706127.